



ZIAUDDIN UNIVERSITY
EXAMINATION BOARD

RESOURCES FOR
“SSC-I PHYSICS”
ZUEB EXAMINATIONS 2021



PREFACE:

The ZUEB examination board acknowledges the serious problems encountered by the schools and colleges in smooth execution of the teaching and learning processes due to sudden and prolonged school closures during the covid-19 spread. The board also recognizes the health, psychological and financial issues encountered by students due to the spread of covid-19.

Considering all these problems and issues the ZUEB Board has developed these resources based on the condensed syllabus 2021 to facilitate students in learning the content through quality resource materials.

The schools and students could download these materials from www.zueb.pk to prepare their students for the high quality and standardized ZUEB examinations 2021.

The materials consist of examination syllabus with specific students learning outcomes per topic, Multiple Choice Questions (MCQs) to assess different thinking levels, Constructed Response Questions (CRQs) with possible answers, Extended Response Questions (ERQs) with possible answers and learning materials.

ACADEMIC UNIT ZUEB:

2. Constructed Response Questions (CRQs)

HOW TO ATTEMPT CRQs:

- Write the answer to each Constructed Response Question/ERQs in the space given below it.
- Use black pen/pencil to write the responses. Do not use glue or pin on the paper.

SECTION B (SHORT ANSWER QUESTIONS)

1. What are physical quantities?

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slight shadow on the right side, suggesting it's resting on a surface. The top edge of the paper is slightly irregular, like a torn piece of paper.

S.NO	CRQ	ANSWER	CL	DL
CHAPTER 2				
1.	What are physical quantities?	There are thousands of physical things and quantities present around us. To understand them and to explain them there must be some scales of measurement so they can be effectively use universally. In 1960 in the international conference, a system of measurements is recommended which is universally acceptable. In this system which is called S.I, seven physical quantities are said to be basic quantities and their scale of measurement are said to be basic units.	K/R	E

Following are these basic quantities and their units.

PHYSICAL QUANTITY	SYM BOL	UNIT	SYM BOL
Length	l	Mete r	m
Mass	m	Kilog ram	Kg
Time	t	Seco nd	S
Current	I	Amp ere	A
Temperature	T	Kelvi n	K
Light intensity	I _v	Cand ela	Cd
Quantity of matter	n	Mole	Mol

S.NO	CRQ	ANSWER	CL	DL
CHAPTER 3				
2.	Define the following: Rest Motion Distance Displacement	<u>REST:</u> “A body is said to be in a state of rest if its position with respect to its surrounding does not change with time.” Examples: A book placed on a table, a fixed pole, buildings etc. <u>MOTION:</u> “A body is said to be in the state of motion if it is continuously changing its position with respect to its surrounding with time.” Examples: A running boy, a flowing stream, moving car, etc. <u>DISTANCE:</u> “It is the length of the actual path of the motion between two points. It may be curve or straight line”.	K/R	E

		It is a scalar quantity. It is usually denoted by 'S' Unit: Unit of distance is meter denoted by (m) <u>DISPLACEMENT</u> “It is the shortest straight line distance between two location directed from one point to the other.” It is a vector quantity. It is usually denoted by 'd'. Unit: Unit of displacement is meter denoted by 'm'.		
3.	Define types of motion?	<u>TYPES OF MOTION</u> All types of motions can be classified as <u>TRANSLATORY (OR) LINEAR MOTION:</u> <i>“If a body is moving on a straight line or curved path but every particle in the body is being displaced by the same amount and not repeating its motion, then it is said to be in linear or translatory motion.”</i> Examples: Motion of a car on a straight road, motion of falling object, motion of motorbike on a circular road etc. <u>ROTATIONAL (OR) CIRCULATORY MOTION:</u> <i>“If a body is moving in a circular path around a fixed point called center and thus repeating its motion with time then this motion is called rotational motion.”</i> Example: Motion of planets around the Sun, electron motion around nucleus, motion of wheels etc. <u>VIBRATORY (OR) OSCILLATORY MOTION:</u> <i>“If a body is moving back and forth or up and down about a fixed point called equilibrium point then this type of motion is known as vibratory or oscillatory motion.”</i> Example: Motion of a pendulum, motion of a swing, motion of bird's feather during flight.	K/R	E
4.	Derive first equation of motion?	<u>FIRST EQUATION OF MOTION</u> Let a body of mass 'm' moving with uniform acceleration "a" starting with initial velocity 'V_i' and attains a final velocity 'V_f' in time 't' then according to the definition of the acceleration, Acceleration = $\frac{\text{change in velocity}}{\text{time}}$	K/A	M

		$a = \frac{\Delta V}{t}$ $a = \frac{V_f - V_i}{t}$ $a t = V_f - V_i$ $V_f = V_i + a t$		
5.	Derive second equation of motion?	SECOND EQUATION OF MOTION Let a body is moving with a uniform acceleration “a” starting with a initial velocity V_i and attains a final velocity V_f in time ‘t’ and covers a distance ‘S’. This distance can be written as, $S = V_{av} \times t \text{ ----- (1)}$ But $V_f + V_i$ $V_{av} = \frac{\quad}{2}$ put in equation (1) $S = \frac{V_f + V_i}{2} \times t \text{ -----(2)}$ from first equation of motion $V_f = V_i + a t$ put in equation (2) $S = \frac{(a t + V_i + V_i)}{2} \times t$ $S = \frac{(a t + 2 V_i)}{2} \times t$ $S = \frac{(a t^2 + 2 t V_i)}{2}$ $S = \frac{a t^2}{2} + \frac{2 t V_i}{2}$ $S = V_i t + \frac{1}{2} a t^2$	K/A	M
6.	A motor cyclist covers 150 m in 10 seconds. Find the speed of the motor cyclist.	15m/s	K/A	E

7.	Find the time taken by sunlight to reach the ground if the distance between the sun and the earth is 1.5×10^8 Km. Velocity of light is 3×10^8 m/s.	8 min 20 seconds	K/A	E
8.	Define and explain Newtons first law of motion?	FIRST LAW OF MOTION Statement: “A body remains at rest or continues to move with constant speed on a straight line unless acted by an unbalance force.” Explanation: This law explains that, if a body is under the influence of several forces and if there is no net force acting on the body then it will keep its state i- e rest or motion with uniform velocity. This law also explains an important property of bodies namely inertia, by which a body tries to maintain its state. Examples: Book placed on a table, motion of a parachute, motion of coin in a viscous fluid.	K/R	M
9.	Define and explain Newtons second law of motion?	SECOND LAW OF MOTION Statement: “It states that whenever an unbalance force is applied on a body it produces acceleration in the body in its own direction, this acceleration is directly proportional to the magnitude of unbalanced force and inversely proportional to the mass of the body.” <i>Mathematical Form:</i> Consider an object of mass “m” on which an unbalance force of magnitude “F” is acting and due to which acceleration “a” is produced in the direction of force. Now according to the first part of the law we can write $a \propto F \text{ ----- (1)}$	K/R	M

		According to the second part of the law $a \propto \frac{1}{m} \quad \text{-----} \quad (2)$ Combining these result $a \propto \frac{F}{m}$ $a = k \frac{F}{m}$ In S.I system constant k = 1 Or $F = m a$																	
10.	Differentiate between mass and weight?	<table><tr><th>S.NO</th><th>MASS</th><th>WEIGHT</th></tr><tr><td>1</td><td>Mass is the quantity of matter in a body and measures the inertia.</td><td>Weight is the force by which earth pulls a body towards its center.</td></tr><tr><td>2</td><td>Mass has no direction.</td><td>Weight is always acts towards the center of earth.</td></tr><tr><td>3</td><td>Mass is a constant quantity and remains same everywhere.</td><td>Weight is different at different distances from center of earth.</td></tr><tr><td>4</td><td>Mass can be measured by a common balance.</td><td>Weight is always measured by spring balance</td></tr></table>	S.NO	MASS	WEIGHT	1	Mass is the quantity of matter in a body and measures the inertia.	Weight is the force by which earth pulls a body towards its center.	2	Mass has no direction.	Weight is always acts towards the center of earth.	3	Mass is a constant quantity and remains same everywhere.	Weight is different at different distances from center of earth.	4	Mass can be measured by a common balance.	Weight is always measured by spring balance	K/R	E
S.NO	MASS	WEIGHT																	
1	Mass is the quantity of matter in a body and measures the inertia.	Weight is the force by which earth pulls a body towards its center.																	
2	Mass has no direction.	Weight is always acts towards the center of earth.																	
3	Mass is a constant quantity and remains same everywhere.	Weight is different at different distances from center of earth.																	
4	Mass can be measured by a common balance.	Weight is always measured by spring balance																	
11.	An object of mass 20 kg is	(60 N)	K/A	M															

	moving with an acceleration of 3 m/s^2 ; find the force acting on it.			
12.	An object of mass 50 kg is moving with an acceleration of 5 m/s^2 ; find the force acting on it.	(250 N)	K/A	E
13.	Define Torque?	DEFINITION “Torque or movement of force is the turning effect of force” FORMULA Torque is the product of the force “F” and force arm “d” Mathematically it can be expressed as $\tau = F \times d$ FORCE ARM Force arm or moment arm is the perpendicular distance between the axis of rotation of the body and line of the action of the force In M.K.S system UNIT the unit of torque is Nm. QUANTITY Torque is a vector quantity POSITIVE TORQUE: Torque is said to be positive if the direction of rotation of the body is anti-clock wise NEGATIVE TORQUE Torque is said to be negative if the direction of rotation of the body is clock wise	K/A	D
14.	Define Equilibrium?	DEFINITION “A body is said to be in equilibrium if it is in rest or move with uniform speed by the influence of force” There are two types of equilibrium STATIC EQUILIBRIUM	K/R	M

		DYNAMIC EQUILIBRIUM STATIC EQUILIBRIUM “A body is said to be in static equilibrium if it is in rest by the influence of force”. EXAMPLE 1) A body lying on the table 2) A body hanging at rest from the ceiling by a string. DYNAMIC EQUILIBRIUM “A body is said to be in dynamic equilibrium if it move with uniform speed by the influence of force”. EXAMPLE 1) Train moving with uniform velocity. 2) Paratrooper falling down with uniform velocity CONDITIONS There are two conditions of equilibrium 1) First condition of equilibrium 2) Second condition of equilibrium FIRST CONDITION OF EQUILIBRIUM The resultant of all the forces acting on a body is zero. OR Algebraic sum of all the forces acting along x-axis and y- axis must equal to zero, mathematically it can be expressed as: $\Sigma F_x = 0$ $\Sigma F_y = 0$ SECOND CONDITION OF EQUILIBRIUM The resultant of all the torque acting on a body is zero Mathematically it can be expressed as: $\Sigma \tau = 0$		
15.	Define centripetal acceleration.	PHYSICAL DEFINITION: “If a body move in a circular path then directions of tangential velocity continuously change. Such an acceleration which produce due to change of direction of tangential velocity called centripetal acceleration.” MATHEMATICAL DEFINITION: It is the ratio of square of the speed to the radius of a circle mathematically it can be expressed as:	K/R	M

		$a_c = \frac{V^2}{r}$ <u>DIRECTION:</u> The direction of centripetal acceleration always towards the centre of a circle		
16.	Define centripetal force and centrifugal force.	CENTRIPETAL FORCE: “Such a force which keeps a body in a circular path called centripetal force.” If a body of mass ‘m’ move in a circular path of radius ‘r’ with uniform speed ‘V’ then body must possess centripetal force and according to Newton’s second law of motion centripetal force can obtained by the formula $F_{cp} = ma_c$ Here $a_c = \frac{V^2}{r}$ we put in above $F_{cp} = \frac{mV^2}{r}$ This is the magnitude of centripetal force direction of centripetal force always towards the centre of a circle. CENTRIFUGAL FORCE: It is the reaction of centripetal force therefore magnitude of centrifugal force is equal to the magnitude of centripetal force but direction of centrifugal force always away from the centre of a circle mathematically it can be express as; $F_{cf} = -F_{cp}$ $F_{cf} = -ma_c$ $F_{cf} = -\frac{mv^2}{r}$	K/R	M
17.	Define work and give its unit.	DEFINITION: “Work is said to be done if a force act on the body and the body displace along the direction of force.” EXPLANATION: If force F and displacement S are in the same direction then work can be obtained by the formula $Work = F S$	K/R	R

		If force F and displacement S makes an angle θ with respect to each other then work can be obtain by the formula Work = $F S \cos \theta$ UNIT: In M.K.S. System unit of work is Joule [J]		
18.	Define Elasticity, Elastic limit, Stress, Strains, Hooks Law, Young's Modulus.	ELASTICITY: When force is applied on a body to change its length, shapes or volume and after removal of this force if a body regains or comes back to its original position then this property is called elasticity. ELASTIC LIMIT: It is the maximum limit of body with in which a body regains its original position after removal of applied force . STRESS: Stress is the amount of reaction force per unit area. Mathematically it can be expressed as: Stress = $\frac{\text{Reaction force}}{\text{area}}$ OR $\sigma = \frac{F}{A}$ In MKS system unit of stress is N/m^2. STRAIN: Strain is the deformation produce by the stress. LONGITUDINAL STRAIN Longitudinal strain is the fractional change in length by the application of stress. Mathematically it can be expressed as $\epsilon = \frac{\text{Change in length}}{\text{Original length}}$ $\epsilon = \frac{\Delta L}{L}$ It has no units. HOOK'S LAW: According to Hook's law, "<u>Within elastic limit stress is directly proportional to strain</u>". Mathematically it can be expressed as: $\therefore$ Stress $\propto$ Strain Stress = k Strain OR	K/R	D

		$\frac{\text{Stress}}{\text{Strain}} = k$ With the help of above equation Hook's Law can also be state as: "Within elastic limit the ratio of stress to the strain always remains unchanged". YOUNG'S MODULUS: Young's Modulus is the ratio of stress to the longitudinal strain. Mathematically it can be expressed as: $Y = \frac{\text{Stress}}{\text{Longitudinal Strain}}$ $Y = \frac{\frac{F}{A}}{\frac{\Delta L}{L}} = \frac{F}{A} \times \frac{L}{\Delta L}$ In MKS system its unit is N/m².													
19.	Give the difference between heat and temperature?	<table><tr><th>HEAT</th><th>TEMPERATURE</th></tr><tr><td>Heat is the form of energy which can transfer from hot body to cold body.</td><td>Temperature measures the degree of hotness or coldness of a body.</td></tr><tr><td>Heat is the total kinetic energy of the molecule of a substance.</td><td>Temperature is the average kinetic energy of the molecule of a substance.</td></tr><tr><td>Physically heat cannot be measured but calculated by the formula $\Delta Q = m c \Delta T$</td><td>Temperature can be measured with the help of thermometer</td></tr><tr><td>In S.I system unit of heat is Joule.</td><td>In S.I system unit of temperature is Kelvin.</td></tr></table>	HEAT	TEMPERATURE	Heat is the form of energy which can transfer from hot body to cold body.	Temperature measures the degree of hotness or coldness of a body.	Heat is the total kinetic energy of the molecule of a substance.	Temperature is the average kinetic energy of the molecule of a substance.	Physically heat cannot be measured but calculated by the formula $\Delta Q = m c \Delta T$	Temperature can be measured with the help of thermometer	In S.I system unit of heat is Joule.	In S.I system unit of temperature is Kelvin.		K/R	M
HEAT	TEMPERATURE														
Heat is the form of energy which can transfer from hot body to cold body.	Temperature measures the degree of hotness or coldness of a body.														
Heat is the total kinetic energy of the molecule of a substance.	Temperature is the average kinetic energy of the molecule of a substance.														
Physically heat cannot be measured but calculated by the formula $\Delta Q = m c \Delta T$	Temperature can be measured with the help of thermometer														
In S.I system unit of heat is Joule.	In S.I system unit of temperature is Kelvin.														
20.	State and explain Boyle's law ,Charles law and pressure law?	BOYLE'S LAW STATEMENT: "At constant temperature and for fix no of molecule volume is inversely proportional to the pressure." EXPLANATION: If 'P' represents pressure and 'V' represents volume of a gas then mathematically		K/R	D										

		Boyle's law can be expressed as: $V \propto \frac{1}{P}$ $V = \frac{K}{P}$ $P V = K$ This is the equation of Boyle's law and with the help of above equation Boyle's law can also be stated as: "At constant temperature and for fix no. of molecule the product of pressure and volume remain constant." CHARLES LAW STATEMENT: "At constant pressure and for fix no. of molecule volume is directly proportional to the temperature." EXPLANATION: If 'V' represent volume of the gas and 'T' represent temperature of a gas then Charles law can be expressed as $V \propto T$ $V = k T$ $\frac{V}{T} = K$ This is the equation of Charles law and with the help of above equation Charles law can also be stated as: "At constant pressure and for fix no. of molecule the ratio of volume to the temperature remain constant" PRESSURE LAW STATEMENT: "At constant volume the pressure of given mass of a gas is directly proportional to its absolute temperature."		
--	--	--	--	--

		$P \propto T$ $P = KT$ $\frac{P}{T} = K$ This is the equation of pressure law and with the help of above equation pressure law can also be stated as: “At constant volume and for fix number of molecules the ratio of pressure to the temperature remains constant.”		
21.	Derive an expression for general gas equation.	General gas equation is a single relation in to which Boyle’s law, Charles law and Avogadro’s law can be combined. According to Boyle’s Law $V \propto \frac{1}{P}$ According to Charles law $V \propto T$ According to Avogadro’s law $V \propto n$ Combining these three laws we get $V \propto \frac{nT}{P}$ $V = \frac{RnT}{P}$ $PV = nRT$ Where R is the constant of proportionality and it is known as universal gas constant or general gas constant. In M.K.S. System its value is 8.313 J/mol K.	K/A	M
22.	Calculate the volume occupied by 5 mole of gas at 27°C if it is subjected to a pressure of 1.0×10^5 N/m² . (gas constant R = 8.13 J mole⁻¹K⁻¹)	[0.12m ³]	U/A	M

23.	If 10 mole of a gas exert a pressure of $20 \times 10^4 \text{ N/m}^2$ when confined in a tank of 40 m^3 capacity at 57°C what would be the pressure of 100 mole of the same gas when confined in a 80 m^3 tank at 477°C .	[$2.27 \times 10^6 \text{ N/m}^2$]	U/A	M
24.	The coefficient of linear thermal expansion of aluminum is $26 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$. An aluminum rod is 2 m long at 25°C . what will be its length at 75°C .	[2.0026 m]	U/A	M
25.	At 16°C the length of an iron rod is 510cm . How long is it at 99°C if the coefficient of linear expansion of iron $\alpha = 12 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$.	[510.508 m]	U/A	M